

- 1. Pedicel of two segments (Fig. 1a) ..... 1 Myrmicinae
- 1. Pedicel of one segment (Fig. 1b) ..... (2)



Fig. 1a



Fig. 1b

- 2. Small (2-3 mm long), dark brown to black, odor of blue cheese when crushed ..... 2 Dolichoderinae
- 2. Not as above ..... (3)

- 3. Dorsum of thorax in profile evenly convex (Fig. 3a) ..... 3 *Camponotus*
- 3. Dorsum of thorax depressed behind level of mesonotum (Fig. 3b) ..... (4)



Fig. 3a



Fig. 3b

- 4. Head and body of uniform color (Fig. 4a) ..... 4 *Formica fusca*
- 4. Head and body bicolored, thorax lighter than head and gaster (Fig. 4b) ..... 5 *Formica rufa*



Fig. 4a



Fig. 4b

- 
- 1. ("Myr") Subfamily Myrmicinae, ants in the genera *Myrmica* or *Leptothorax*.
  - 2. ("Dol") Subfamily Dolichoderinae. Only one species, *Tapinoma sessile*.
  - 3. ("Camp") Subfamily Formicinae, ants in the genus *Camponotus*.
  - 4. ("fusca") Subfamily Formicinae, genera *Formica*, species group *fusca* as well as species group *lasioides*.
  - 5. ("rufa") Subfamily Formicinae, genera *Formica*, species in the species group *rufa* as well as the species group *sanguinea*.